

Examples for kinetic evaluations using mkin

Johannes Ranke

Eurofins Regulatory AG
Weidenweg 15, CH-4310 Rheinfelden, Switzerland

and

University of Bremen

February 18, 2013

Contents

1 Kinetic evaluations for parent compounds	1
1.1 Laboratory Data L1	1
1.2 Laboratory Data L2	6
1.3 Laboratory Data L3	14
1.4 Laboratory Data L4	19

Key words: Kinetics, FOCUS, nonlinear optimisation

1 Kinetic evaluations for parent compounds

These examples are also evaluated in a parallel vignette of the **kinfit** package ([Ranke, 2012](#)). The datasets are from Appendix 3, of the FOCUS kinetics report ([FOCUS Work Group on Degradation Kinetics, 2006, 2011](#)).

1.1 Laboratory Data L1

The following code defines example dataset L1 from the FOCUS kinetics report, p. 284

```
R> library("mkin")
R> FOCUS_2006_L1 = data.frame(
+   t = rep(c(0, 1, 2, 3, 5, 7, 14, 21, 30), each = 2),
+   parent = c(88.3, 91.4, 85.6, 84.5, 78.9, 77.6,
+             72.0, 71.9, 50.3, 59.4, 47.0, 45.1,
+             27.7, 27.3, 10.0, 10.4, 2.9, 4.0))
R> FOCUS_2006_L1_mkin <- mkin_wide_to_long(FOCUS_2006_L1)
```

The next step is to set up the models used for the kinetic analysis. Note that the model definitions contain the names of the observed variables in the data. In this case, there is only one variable called `parent`.

```
R> SFO <- mkinmod(parent = list(type = "SFO"))
R> FOMC <- mkinmod(parent = list(type = "FOMC"))
R> DFOP <- mkinmod(parent = list(type = "DFOP"))
```

The three models cover the first assumption of simple first order (SFO), the case of declining rate constant over time (FOMC) and the case of two different phases of the kinetics (DFOP). For a more detailed discussion of the models, please see the FOCUS kinetics report.

The following two lines fit the model and produce the summary report of the model fit. This covers the numerical analysis given in the FOCUS report.

```

R> m.L1.SFO <- mkinfit(SFO, FOCUS_2006_L1_mkin, quiet=TRUE)
R> summary(m.L1.SFO)

mkin version: 0.9.11
R version: 2.15.2
Date of fit: Mon Feb 18 00:33:32 2013
Date of summary: Mon Feb 18 00:33:32 2013

Equations:
[1] d_parent = - k_parent_sink * parent

Starting values for optimised parameters:
      initial   type transformed
parent_0       100.0 state 100.000000
k_parent_sink     0.1 deparm -2.302585

Fixed parameter values:
None

Optimised, transformed parameters:
      Estimate Std. Error
parent_0       92.471    1.368
k_parent_sink   -2.347    0.041

Backtransformed parameters:
      Estimate
parent_0       92.471
k_parent_sink   0.096

Residual standard error: 2.948 on 16 degrees of freedom

Chi2 error levels in percent:
      err.min n.optim df
All data     3.424      2   7
parent       3.424      2   7

Estimated disappearance times:
      DT50   DT90
parent 7.249 24.08

Estimated formation fractions:
      ff
parent_sink  1

Parameter correlation:
      parent_0 k_parent_sink
parent_0       1.0000      0.6248
k_parent_sink   0.6248      1.0000


```

Data:

time	variable	observed	predicted	residual
0	parent	88.3	92.471	-4.1710
0	parent	91.4	92.471	-1.0710
1	parent	85.6	84.039	1.5610
1	parent	84.5	84.039	0.4610
2	parent	78.9	76.376	2.5241
2	parent	77.6	76.376	1.2241
3	parent	72.0	69.412	2.5884
3	parent	71.9	69.412	2.4884
5	parent	50.3	57.330	-7.0301
5	parent	59.4	57.330	2.0699
7	parent	47.0	47.352	-0.3515
7	parent	45.1	47.352	-2.2515
14	parent	27.7	24.247	3.4527
14	parent	27.3	24.247	3.0527
21	parent	10.0	12.416	-2.4163
21	parent	10.4	12.416	-2.0163
30	parent	2.9	5.251	-2.3513
30	parent	4.0	5.251	-1.2513

A plot of the fit is obtained with the plot function for mkinfit objects.


```
R> plot(m.L1.SF0)
```



The residual plot can be easily obtained by

```
R> mkinresplot(m.L1.SFO, ylab = "Observed", xlab = "Time")
```

Residuals of mkin fit



For comparison, the FOMC model is fitted as well, and the χ^2 error level is checked.

```
R> m.L1.FOMC <- mkinfit(FOMC, FOCUS_2006_L1_mkin, quiet=TRUE)
R> s.m.L1.FOMC <- summary(m.L1.FOMC)
R> s.m.L1.FOMC$errmin
      err.min n.optim df
All data 0.03618911      3   6
parent    0.03618911      3   6
```

Due to the higher number of parameters, and the lower number of degrees of freedom of the fit, the χ^2 error level is actually higher for the FOMC model (3.6%) than for the SFO model (3.4%).

1.2 Laboratory Data L2

The following code defines example dataset L2 from the FOCUS kinetics report, p. 287

```
R> FOCUS_2006_L2 = data.frame(  
+   t = rep(c(0, 1, 3, 7, 14, 28), each = 2),  
+   parent = c(96.1, 91.8, 41.4, 38.7,  
+             19.3, 22.3, 4.6, 4.6,  
+             2.6, 1.2, 0.3, 0.6))  
R> FOCUS_2006_L2_mkin <- mkin_wide_to_long(FOCUS_2006_L2)
```

Again, the SFO model is fitted and a summary is obtained.

```
R> m.L2.SFO <- mkinfit(SFO, FOCUS_2006_L2_mkin, quiet=TRUE)  
R> summary(m.L2.SFO)
```

```
mkin version: 0.9.11  
R version: 2.15.2  
Date of fit: Mon Feb 18 00:33:32 2013  
Date of summary: Mon Feb 18 00:33:32 2013
```

Equations:

```
[1] d_parent = - k_parent_sink * parent
```

Starting values for optimised parameters:

	initial	type	transformed
parent_0	100.0	state	100.000000
k_parent_sink	0.1	deparm	-2.302585

Fixed parameter values:

None

Optimised, transformed parameters:

	Estimate	Std. Error
parent_0	91.4656	3.807
k_parent_sink	-0.4112	0.107

Backtransformed parameters:

	Estimate
parent_0	91.466
k_parent_sink	0.663

Residual standard error: 5.51 on 10 degrees of freedom

Chi2 error levels in percent:

	err.min	n.optim	df
All data	14.38	2	4
parent	14.38	2	4

Estimated disappearance times:

	DT50	DT90
parent	1.046	3.474

Estimated formation fractions:

	ff
parent_sink	1

Parameter correlation:


	parent_0	k_parent_sink
parent_0	1.0000	0.4295
k_parent_sink	0.4295	1.0000

Data:

time	variable	observed	predicted	residual
0	parent	96.1	91.4656079103	4.6344
0	parent	91.8	91.4656079103	0.3344
1	parent	41.4	47.1395280371	-5.7395
1	parent	38.7	47.1395280371	-8.4395
3	parent	19.3	12.5210295280	6.7790
3	parent	22.3	12.5210295280	9.7790
7	parent	4.6	0.8833842647	3.7166
7	parent	4.6	0.8833842647	3.7166
14	parent	2.6	0.0085318162	2.5915
14	parent	1.2	0.0085318162	1.1915
28	parent	0.3	0.0000007958	0.3000
28	parent	0.6	0.0000007958	0.6000

The χ^2 error level of 14% suggests that the model does not fit very well. This is also obvious from the plots of the fit and the residuals.


R> plot(m.L2.SF0)



In the FOCUS kinetics report, it is stated that there is no apparent systematic error observed from the residual plot up to the measured DT90 (approximately at day 5), and there is an underestimation beyond that point.

```
R> mkinresplot(m.L2.SF0, ylab = "Observed", xlab = "Time")
```


Residuals of mkin fit



We may add that it is difficult to judge the random nature of the residuals just from the three samplings at days 0, 1 and 3. Also, it is not clear why a consistent underestimation after the approximate DT90 should be irrelevant.

For comparison, the FOMC model is fitted as well, and the χ^2 error level is checked.


```
R> m.L2.FOMC <- mkinfit(FOMC, FOCUS_2006_L2_mkin, quiet=TRUE)
R> plot(m.L2.FOMC)
R> s.m.L2.FOMC <- summary(m.L2.FOMC)
R> s.m.L2.FOMC$errmin
      err.min n.optim df
All data 0.06204245      3   3
parent    0.06204245      3   3
```



The error level at which the χ^2 test passes is much lower in this case. Therefore, the FOMC model provides a better description of the data, as less experimental error has to be assumed in order to explain the data.

Fitting the four parameter DFOP model does not further reduce the χ^2 error level.

```
R> m.L2.DFOP <- mkinfit(DFOP, FOCUS_2006_L2_mkin, quiet=TRUE)
R> plot(m.L2.DFOP)
```



Here, the default starting parameters for the DFOP model obviously do not lead to a reasonable solution. Therefore the fit is repeated with different starting parameters.

```
R> m.L2.DFOP <- mkinfit(DFOP, FOCUS_2006_L2_mkin,
+   parms.ini = c(k1 = 1, k2 = 0.01, g = 0.8),
+   quiet=TRUE)
R> plot(m.L2.DFOP)
R> summary(m.L2.DFOP)

mkin version: 0.9.11
R version: 2.15.2
Date of fit: Mon Feb 18 00:33:33 2013
Date of summary: Mon Feb 18 00:33:33 2013
```

Equations:

```
[1] d_parent = - ((k1 * g * exp(-k1 * time) + k2 * (1 - g) * exp(-k2 * time)) / (g * exp
```

Starting values for optimised parameters:

	initial	type	transformed
parent_0	1e+02	state	100.0000000
k1	1e+00	deparm	0.0000000
k2	1e-02	deparm	-4.6051702
g	8e-01	deparm	0.9802581

Fixed parameter values:

None

Optimised, transformed parameters:

	Estimate	Std. Error
parent_0	93.9500	NA
k1	4.9589	NA
k2	-1.0880	NA
g	-0.2821	NA

Backtransformed parameters:

	Estimate
parent_0	93.950
k1	142.434
k2	0.337
g	0.402

Residual standard error: 1.732 on 8 degrees of freedom

Chi2 error levels in percent:

	err.min	n.optim	df
All data	2.529	4	2
parent	2.529	4	2

Estimated disappearance times:

	DT50	DT90
parent	NA	NA

Estimated formation fractions:

```
[1] ff  
<0 Zeilen> (oder row.names mit Länge 0)
```

Data:

	time	variable	observed	predicted	residual
0	parent		96.1	93.950000	2.1500
0	parent		91.8	93.950000	-2.1500
1	parent		41.4	40.143423	1.2566
1	parent		38.7	40.143423	-1.4434


```

3  parent      19.3 20.464500 -1.1645
3  parent      22.3 20.464500  1.8355
7  parent      4.6  5.318322 -0.7183
7  parent      4.6  5.318322 -0.7183
14 parent      2.6   0.503070  2.0969
14 parent      1.2   0.503070  0.6969
28 parent      0.3   0.004501  0.2955
28 parent      0.6   0.004501  0.5955

R> s.m.L2.DFOP <- summary(m.L2.DFOP)
R> s.m.L2.DFOP$errmin

      err.min n.optim df
All data 0.02528763      4  2
parent    0.02528763      4  2

```



Therefore, the FOMC model is clearly the best-fit model based on the χ^2 error level criterion.

1.3 Laboratory Data L3

The following code defines example dataset L3 from the FOCUS kinetics report, p. 290

```
R> FOCUS_2006_L3 = data.frame(
+   t = c(0, 3, 7, 14, 30, 60, 91, 120),
+   parent = c(97.8, 60, 51, 43, 35, 22, 15, 12))
R> FOCUS_2006_L3_mkin <- mkin_wide_to_long(FOCUS_2006_L3)
```

SFO model, summary and plot:

```
R> m.L3.SFO <- mkinfit(SFO, FOCUS_2006_L3_mkin, quiet=TRUE)
R> summary(m.L3.SFO)

mkin version: 0.9.11
R version: 2.15.2
Date of fit: Mon Feb 18 00:33:34 2013
Date of summary: Mon Feb 18 00:33:34 2013

Equations:
[1] d_parent = - k_parent_sink * parent

Starting values for optimised parameters:
      initial   type transformed
parent_0       100.0 state 100.000000
k_parent_sink     0.1 deparm -2.302585

Fixed parameter values:
None

Optimised, transformed parameters:
      Estimate Std. Error
parent_0    74.873    8.458
k_parent_sink -3.678    0.326

Backtransformed parameters:
      Estimate
parent_0    74.873
k_parent_sink 0.025

Residual standard error: 12.91 on 6 degrees of freedom

Chi2 error levels in percent:
      err.min n.optim df
All data    21.24      2   6
parent      21.24      2   6

Estimated disappearance times:
      DT50   DT90
parent 27.43 91.12

Estimated formation fractions:
      ff
parent_sink  1


Parameter correlation:
      parent_0 k_parent_sink
parent_0      1.0000      0.5484
```

```
k_parent_sink 0.5484      1.0000
```

Data:

time	variable	observed	predicted	residual
0	parent	97.8	74.873	22.92734
3	parent	60.0	69.407	-9.40654
7	parent	51.0	62.734	-11.73403
14	parent	43.0	52.563	-9.56336
30	parent	35.0	35.083	-0.08281
60	parent	22.0	16.439	5.56137
91	parent	15.0	7.510	7.48961
120	parent	12.0	3.609	8.39083

```
R> plot(m.L3.SFO)
```



The χ^2 error level of 22% as well as the plot suggest that the model does not fit very well.

The FOMC model performs better:

```
R> m.L3.FOMC <- mkinfit(FOMC, FOCUS_2006_L3_mkin, quiet=TRUE)
R> plot(m.L3.FOMC)
R> s.m.L3.FOMC <- summary(m.L3.FOMC)
R> s.m.L3.FOMC$errmin


      err.min n.optim df
All data 0.07321867      3   5
parent    0.07321867      3   5

R> endpoints(m.L3.FOMC)

$distimes
      DT50      DT90
parent 7.729478 431.2428

$ff
logical(0)


$SFORB
logical(0)
```



The error level at which the χ^2 test passes is 7% in this case.

Fitting the four parameter DFOP model further reduces the χ^2 error level considerably:

```
R> m.L3.DFOP <- mkinfit(DFOP, FOCUS_2006_L3_mkin, quiet=TRUE)
R> plot(m.L3.DFOP)
R> s.m.L3.DFOP <- summary(m.L3.DFOP)
R> s.m.L3.DFOP$errmin
      err.min n.optim df
All data 0.02223992      4   4
parent    0.02223992      4   4
```



Therefore, the DFOP model is the best-fit model based on the χ^2 error level criterion for laboratory data L3.

1.4 Laboratory Data L4

The following code defines example dataset L4 from the FOCUS kinetics report, p. 293

```
R> FOCUS_2006_L4 = data.frame(
+   t = c(0, 3, 7, 14, 30, 60, 91, 120),
+   parent = c(96.6, 96.3, 94.3, 88.8, 74.9, 59.9, 53.5, 49.0))
R> FOCUS_2006_L4_mkin <- mkin_wide_to_long(FOCUS_2006_L4)
```

SFO model, summary and plot:

```
R> m.L4.SFO <- mkinfit(SFO, FOCUS_2006_L4_mkin, quiet=TRUE)
R> summary(m.L4.SFO)

mkin version: 0.9.11
R version: 2.15.2
Date of fit: Mon Feb 18 00:33:34 2013
Date of summary: Mon Feb 18 00:33:34 2013

Equations:
[1] d_parent = - k_parent_sink * parent

Starting values for optimised parameters:
      initial   type transformed
parent_0       100.0 state 100.000000
k_parent_sink     0.1 deparm -2.302585

Fixed parameter values:
None

Optimised, transformed parameters:
      Estimate Std. Error
parent_0      96.44     1.949
k_parent_sink -5.03     0.080

Backtransformed parameters:
      Estimate
parent_0      96.442
k_parent_sink  0.007

Residual standard error: 3.651 on 6 degrees of freedom

Chi2 error levels in percent:
      err.min n.optim df
All data    3.288      2   6
parent      3.288      2   6

Estimated disappearance times:
      DT50 DT90
parent    106  352

Estimated formation fractions:
      ff
parent_sink  1


Parameter correlation:
      parent_0 k_parent_sink
parent_0      1.0000     0.5865
```

```
k_parent_sink    0.5865          1.0000
```

Data:

time	variable	observed	predicted	residual
0	parent	96.6	96.44	0.1585
3	parent	96.3	94.57	1.7324
7	parent	94.3	92.13	2.1744
14	parent	88.8	88.00	0.7972
30	parent	74.9	79.26	-4.3589
60	parent	59.9	65.14	-5.2376
91	parent	53.5	53.18	0.3167
120	parent	49.0	43.99	5.0054

```
R> plot(m.L4.SFO)
```




The χ^2 error level of 3.3% as well as the plot suggest that the model fits very well.

The FOMC model for comparison

```
R> m.L4.FOMC <- mkinfit(FOMC, FOCUS_2006_L4_mkin, quiet=TRUE)
R> plot(m.L4.FOMC)
R> s.m.L4.FOMC <- summary(m.L4.FOMC)
R> s.m.L4.FOMC$errmin
```

	err.min	n.optim	df
All data	0.02027643	3	5
parent	0.02027643	3	5



The error level at which the χ^2 test passes is slightly lower for the FOMC model. However, the difference appears negligible.

References

- FOCUS Work Group on Degradation Kinetics. *Guidance Document on Estimating Persistence and Degradation Kinetics from Environmental Fate Studies on Pesticides in EU Registration. Report of the FOCUS Work Group on Degradation Kinetics*, 2006. URL <http://focus.jrc.ec.europa.eu/dk>. EC Document Reference Sanco/10058/2005 version 2.0.
- FOCUS Work Group on Degradation Kinetics. *Generic guidance for estimating persistence and degradation kinetics from environmental fate studies on pesticides in EU registration*, 1.0 edition, November 2011. URL <http://focus.jrc.ec.europa.eu/dk>.
- Johannes Ranke. *kinfit: Routines for fitting simple kinetic models to chemical degradation data*, 2012. URL <http://CRAN.R-project.org>.